

FACTORIZING 3-FOLD FLIPS AND DIVISORIAL CONTRACTIONS TO CURVES

JUNGKAI A. CHEN AND CHRISTOPHER D. HACON

ABSTRACT. We show that 3-fold terminal flips and divisorial contractions to a curve may be factored by a sequence of weighted blow-ups, flops, blow-downs to a locally complete intersection curve in a smooth 3-fold or divisorial contractions to a point.

1. INTRODUCTION

Flips, flops and divisorial contractions are the “elementary” birational maps of the minimal model program. Divisorial contractions are the higher dimensional analog of (the inverse of) blowing up a smooth point on a surface. They are morphisms contracting a divisor covered by K_X -negative curves. Flips are instead a new operation which consists of a “surgery” in codimension ≥ 2 which replaces certain K_X -negative curves by some K_X -positive curves. Flops are also surgeries in codimension ≥ 2 but they replace certain K_X -trivial curves by some new K_X -trivial curves.

These operations appear as follows. Given a smooth projective variety of general type over $\mathbb{C}$, then a minimal model of X may be constructed via a finite sequence of flips and divisorial contractions (cf. [1]). Minimal models are not unique, but it is known that any two minimal models are connected by a finite sequence of flops (cf. [16]).

Not surprisingly, the geometry of divisorial contractions is better understood than that of flips and flops. In fact the proof of the existence of flips and flops (cf. [1], [7]) is very abstract and gives little insight to their geometry.

In dimension 3 these operations are reasonably well understood. Kollár has shown [23] that if $X \dashrightarrow X^+$ is a flop (of terminal 3-folds), then X and X^+ have analytically isomorphic singularities. In his seminal paper [34], Mori gives a precise understanding of the geometry of all flips with irreducible flipping curve (extremal neighborhoods).

The first author was partially supported by TIMS, NCTS/TPE and National Science Council of Taiwan. The second author was supported by NSF grant 0757897. We are indebted to Hayakawa, Kollár and Mori for many useful discussions especially to Mori who read our preliminary version and gave various comments and corrections. Some of this work was done during a visit of the first author to the University of Utah. The first author would like to thank the University of Utah for its hospitality.

It should also be noted that 3-fold divisorial contractions may be subdivided into two classes: those that map the divisor to a curve and those that map the divisor to a point. Divisorial contractions to a point are classified in [4], [14], [18], [19], [20], [21] and [31]. There are also several partial results on divisorial contractions to a curve cf. [5], [37], [38], [39], [40] and [41].

The purpose of this paper is to show that in dimension 3, we can factor flips and divisorial contractions to curves via the simpler operations given by flops, blow-downs to LCI curves (i.e. $C \subset Y$ a local complete intersection curve in a smooth variety) and divisorial contractions to points. More precisely we prove

Theorem 1.1. *Let $g : X \rightarrow W$ be a flipping contraction and $\phi : X \dashrightarrow X^+$ be the corresponding flip, then ϕ can be factored as*

$$X = X_0 \xrightarrow{f_0} X_1 \dashrightarrow \dots \dashrightarrow X_n \xrightarrow{f_n} X^+,$$

such that each f_i is the inverse of a w -morphism, a flop, a blow-down to a LCI curve or a divisorial contraction to a point.

Let $g : X \rightarrow W$ be a divisorial contraction to a curve, then g can be factored as

$$X = X_0 \xrightarrow{f_0} X_1 \dashrightarrow \dots \dashrightarrow X_n \xrightarrow{f_n} W,$$

such that each f_i is the inverse of a w -morphism, a flop, a blow-down to a LCI curve or a divisorial contraction to a point.

We remark that the proof of the above theorem follows by the classification results of [34] and the results of [8] and [9]. One should also note that this kind of approach is not entirely new. See for example [13] and [11].

As a final motivation for (1.1), we would like to mention a possible application of these ideas to characteristic $p > 0$. In this context, the existence of divisorial contractions (in the big case) is known by work of Keel [22]. The existence of flips is instead only known in some special cases [13] and [24]. Theorem 1.1 suggests that it may be possible to construct flips in characteristic $p > 0$ via a sequence of divisorial contractions (and extractions) and flops (which are hopefully easier to construct than flips).

2. PRELIMINARIES

We work over the field of complex numbers $\mathbb{C}$. *Numerical* (resp. *linear*, $\mathbb{Q}$ -*linear*) *equivalence* of divisors is denoted by $\equiv$ (resp. $\sim$, $\sim_{\mathbb{Q}}$), if $f : X \rightarrow Y$ is a projective morphism, then $\equiv_Y$ denotes *f-numerical* (also known as numerical over Y) *equivalence* and $\sim_Y$ (resp. $\sim_{\mathbb{Q},Y}$) denote *linear equivalence* (resp. $\mathbb{Q}$ -*linear equivalence*) *over* Y . So, for example, $D \sim_{\mathbb{Q},Y} D'$ if and only if there are rational numbers $q_i \in \mathbb{Q}$, rational functions $f_i \in \mathbb{C}(X)$ and a $\mathbb{Q}$ -Cartier divisor L on Y such that

$D - D' = \sum q_i(f_i) + f^*L$. If X is a normal variety, then K_X denotes a canonical divisor on X . A *log pair* (X, B) is given by a normal variety X and a $\mathbb{Q}$ -divisor B such that $K_X + B$ is $\mathbb{Q}$ -Cartier. A *log resolution* of a log pair (X, B) is a proper birational morphism $\nu : X' \rightarrow X$ such that $\text{Exc}(\nu) \cup \nu_*^{-1}B$ is a simple normal crossings divisor. If (X, B) is a log pair and $\nu : X' \rightarrow X$ is a log resolution of (X, B) , then we write $K_{X'} + \nu_*^{-1}B = \nu^*(K_X + B) + \sum a_i E_i$ where $\nu_* K_{X'} = K_X$. The rational numbers $a_{E_i}(X, B) := a_i$ are the *discrepancies* of (X, B) . The log pair (X, B) is *terminal* (resp. *canonical* or *plt*) if $a_E(X, B) > 0$ (resp. $a_E(X, B) \geq 0$ or $a_E(X, B) > -1$) for all exceptional divisors E over X . The log pair $(X, B = \sum b_i B_i)$ is *klt* if it is plt and $b_i < 1$ for all i . By inversion of adjunction, it is known that if $(X, S + B)$ is a log pair with $\lfloor S + B \rfloor = S$, then $(X, S + B)$ is plt (resp. canonical or terminal) if and only if (S, B_S) is klt (resp. canonical or terminal) where $(K_X + S + B)|_S = K_S + B_S$.

2.1. Terminal threefold singularities. Here we briefly recall several facts about terminal singularities in dimension 3. It is well known that terminal surfaces are smooth and hence terminal threefolds have isolated singularities. Let $P \in X$ be the germ of a terminal singularity, then X has rational singularities. The *index* of $P \in X$ is the smallest integer $r > 0$ such that rK_X is Cartier on a neighborhood of P . If $P \in X$ is a point of index r and D is any Weil divisor on X , then rD is also Cartier on a neighborhood of P . The canonical cover $\pi : (P^\# \in X^\#) \rightarrow (P \in X)$ is a cyclic cover of degree r such that $P^\#$ is an isolated *cDV* point of index 1 (cf. [36, 3.1]). The analytic germ $(P^\# \in X^\#)$ can be embedded into $\mathbb{C}^4$ as a hypersurface $\varphi = 0$. The singularity $(P \in X)$ can thus be realized as a cyclic group μ_r -quotient of a hypersurface singularity. Note that there exists a semi-invariant coordinate system $y_1, \dots, y_4$ near $(P^\# \in X^\#)$. These singularities are classified in [33] (see also [34, §1a] for a brief survey).

The weights of the coordinates $y_1, \dots, y_4$ are usually denoted by $\frac{1}{r}(a_1, a_2, a_3, a_4)$ [34, pg. 243]. Two weights $\frac{1}{r}(a_1, a_2, a_3, a_4)$ and $\frac{1}{r}(b_1, b_2, b_3, b_4)$ are said to be equivalent if there is an integer λ relatively prime to r , such that $b_i \equiv a_i \lambda \pmod{r}$ for all i . Recall that for any terminal singularity $P \in X$ there is a deformation producing $k \geq 1$ terminal cyclic quotient singularities $P_1, \dots, P_k$. The number $k = aw(P \in X)$ is called the *axial weight* of $(P \in X)$. We let

$$aw(X) = \sum_{P \in \text{Sing}(X)} aw(P \in X).$$

Note that the axial weight coincides with the *axial multiplicity* defined in [34, Definition-Corollary 1a.5.iii] except in the case $cAx/4$. We may assume that the singularities P_i have type $\frac{1}{r_i}(1, -1, b_i)$ where $0 < b_i \leq r/2$. The collection $\{P_1, \dots, P_k\}$ is known as the *basket of singularities*

of (X, P) and it can be written as

$$\mathcal{B}(P \in X) = \{n_i \times (b_i, r_i) | i \in I, n_i \in \mathbb{Z}^+\},$$

where n_i denotes the number of times that a point P_i representing a singularity $\frac{1}{r_i}(1, -1, b_i)$ appears.

To a given variety X , we can associate the basket of singularities $\mathcal{B} = \cup_{P \in \text{Sing}(X)} \mathcal{B}(P \in X)$. We can define the following invariants corresponding to a singularity

$$\begin{aligned} \sigma(P \in X) &= \sum_{i=1}^{aw(P \in X)} b(P_i), & \sigma(X) &= \sum_{P \in \text{Sing}(X)} \sigma(P \in X), \\ \Xi(P \in X) &= \sum_{i=1}^{aw(P \in X)} r(P_i), & \Xi(X) &= \sum_{P \in \text{Sing}(X)} \Xi(P \in X). \end{aligned}$$

Remark 2.1. The basket $\mathcal{B}(P \in X)$ is uniquely determined by the singularity $(P \in X)$. We have

type	G. E.	aw	basket	σ	Ξ
cA/r	A_{kr-1}	k	$k \times (b, r)$	kb	kr
$cAx/2$	D_{k+2}	2	$2 \times (1, 2)$	2	4
$cAx/4$	D_{2k+1}	k	$\{(1, 4), (k-1) \times (1, 2)\}$	k	$2k+2$
$cD/2$	D_{2k}	k	$k \times (1, 2)$	k	$2k$
$cD/3$	E_6	2	$2 \times (1, 3)$	2	6
$cE/2$	E_7	3	$3 \times (1, 2)$	2	6

where the column G. E. lists the singularities for general elephant (i.e. a general element) $E_X \in |-K_X|$.

2.2. Extremal neighborhoods. Recall that (cf. [30, §1]) an *extremal neighborhood* $f : X \supset C \rightarrow Y \ni Q$ is a proper bimeromorphic morphism of complex 3 dimensional spaces $f : X \rightarrow Y$ such that

- (1) X is terminal;
- (2) Y is normal with distinguished point $Q \in Y$;
- (3) $f^{-1}(Q)$ consists of a single irreducible curve $C \subset X$;
- (4) $K_X \cdot C < 0$.

If instead of the condition $K_X \cdot C < 0$, we have the condition $K_X \cdot C = 0$, then we will say that f is a *K-trivial extremal neighborhood*. An *isolated extremal neighborhood* is an extremal neighborhood $f : X \supset C \rightarrow Y \ni Q$ such that $\dim \text{Exc}(f) = 1$. In this case $\text{Exc}(f) = C$ and K_Y is not $\mathbb{Q}$ -Cartier.

An extremal neighborhood is *divisorial* if $\dim \text{Exc}(f) = 2$. In this case, $\text{Exc}(f)$ is an irreducible divisor, K_Y is $\mathbb{Q}$ -Cartier and Y is terminal.

By [34, pg. 151, 2.3.2],

$$(1) \quad K_X \cdot C = -1 + \sum_{P \in C} w_P(0),$$

where

$$w_P(0) = \begin{cases} 0 & \text{if } P \text{ is of index } 1, \\ 0 \leq \frac{c}{r} \leq \frac{r-1}{r} & \text{if } P \text{ is of index } r > 1. \end{cases}$$

The exact value of $w_P(0)$ is given in [34, pg. 175, 4.9].

Extremal neighborhoods are classified in [34], see also [30] and [35]. We have the following fundamental result also known as Reid's general elephant conjecture (cf. [34]).

Theorem 2.2. *Let $f : X \supset C \rightarrow Y \ni Q$ be an extremal neighborhood and $E_X \in |-K_X|$ and $E_Y \in |-K_Y|$ be general elements. Then E_X and E_Y are normal and have only Du Val singularities.*

2.3. Extremal contractions. A *flipping contraction* (resp. *flopping contraction*) is a projective birational morphism $f : X \rightarrow Y$ from a terminal 3-fold to a normal variety such that $\dim \text{Exc}(f) = 1$, $\rho(X/Y) = 1$, $f_*\mathcal{O}_X = \mathcal{O}_Y$ and $-K_X$ is f -ample (resp. K_X is f -trivial). A *flipping curve* (resp. *flopping curve*) is a curve $C \subset \text{Exc}(f)$. The *flip* (resp. *flop*) of a flipping contraction (resp. a flopping contraction) $f : X \rightarrow Y$ is a birational morphism $f^+ : X^+ \rightarrow Y$ where X^+ is a terminal 3-fold, $\dim \text{Exc}(f^+) = 1$, $\rho(X^+/Y) = 1$, $f^+_*\mathcal{O}_{X^+} = \mathcal{O}_Y$ and K_{X^+} is f -ample (resp. f -trivial). A *flipped curve* (resp. a *flopped curve*) is a curve $C \subset \text{Exc}(f^+)$. One defines the flip (resp. flop) of an extremal neighborhood (resp. K -trivial extremal neighborhood) analogously.

A *divisorial contraction* is a birational morphism $f : X \rightarrow Y$ from a terminal 3-fold to a normal variety such that $\dim \text{Exc}(f) = 2$, $\rho(X/Y) = 1$, $f_*\mathcal{O}_X = \mathcal{O}_Y$ and $-K_X$ is f -ample. It is known that the contracted divisor $F \subset X$ is irreducible and Y is terminal.

We will need the following results.

Theorem 2.3. *Let $f : X \supset C \rightarrow Y \ni P$ be a flipping (resp. flopping contraction). Then the flip (resp. flop) $f^+ : X^+ \rightarrow Y$ exists. Moreover, in the analytic category, $\varphi : X \dashrightarrow X^+$ factors via a finite sequence of flips (resp. flops) $\varphi_i : X_i \dashrightarrow X_i^+ = X_{i+1}$ with irreducible flipping (resp. flopping) curves $f_i : X_i \rightarrow Y_i$ over Y such that $\text{Exc}(\varphi_i) = C_i \cong \mathbb{P}^1$ and $\text{Exc}(\varphi_i^+) = C_i^+ \cong \mathbb{P}^1$.*

Proof. See [34] and [15]. □

Remark 2.4. Notice that the arguments of [15] also imply that if $f : X \supset C \rightarrow Y \ni P$ is a contraction of a divisor to a curve and $C = \cup C_j$ is not irreducible, then (in the analytic category) f factors through an extremal neighborhood.

Remark 2.5. It is expected that if X is a terminal 3-fold and $f : X \rightarrow Y$ is the germ of a flipping or divisorial contraction to a curve and if $E_X \in |-K_X|$ is general, then the divisor E_X is normal with Du Val singularities.

Theorem 2.6. *Let X be a terminal Gorenstein 3-fold, $f : X \rightarrow Y$ an extremal contraction. Then f is divisorial and if it contracts the divisor to a curve C , then X has only A_n singularities, $C \subset Y$ is LCI (i.e. Y is smooth near C and C is a local complete intersection in Y) and X is the blow up of Y along C . If f contracts the divisor to a point $P \in Y$, then $f : X \rightarrow Y$ is classified (it belongs to one of the four possibilities listed in [5]).*

Proof. See [5]. □

We will need the following easy lemmas.

Lemma 2.7. *Let $P \in X$ be a terminal 3-fold singularity of index $r > 1$ and $f : Y \supset F \rightarrow X \ni P$ be an extremal divisorial contraction of discrepancy $1/r$. Let $G_X \in |-K_X|$ be a normal divisor with only Du Val singularities and G_Y its strict transform on Y , then*

- (1) $P \in \text{Supp}(G_X)$,
- (2) $f^*(K_X + G_X) = K_Y + G_Y$,
- (3) $f|_{G_Y} : G_Y \rightarrow G_X$ is not an isomorphism, and
- (4) $f|_{G_Y}$ is dominated by the minimal resolution $\tilde{G}_X \rightarrow G_X$.

Proof. We have that $f^*(K_X + G_X) = K_Y + G_Y + tF$. Since $-K_X$ is not Cartier at P , we have that $P \in \text{Supp}(G_X)$. It follows that $t \geq 0$. Since G_X is canonical, by inversion of adjunction, (X, G_X) is canonical. Therefore, $t \leq 0$ and hence $t = 0$. Since F is a $\mathbb{Q}$ -divisor, so is $G_Y \cap F \neq \emptyset$. It follows that $f|_{G_Y}$ is not an isomorphism. Since G_X is Du Val and $K_{G_Y} = (f|_{G_Y})^*K_{G_X}$, $f|_{G_Y}$ is dominated by the minimal resolution $\tilde{G}_X \rightarrow G_X$. □

Lemma 2.8. *Let X be a terminal analytic 3-fold and $G_X \in |-K_X|$ be a normal divisor with only Du Val singularities. If $E_X \in |-K_X|$ is general, then E_X is a normal divisor with only Du Val singularities and $\rho(\tilde{E}_X/E_X) \leq \rho(\tilde{G}_X/G_X)$ where $\tilde{E}_X$ and $\tilde{G}_X$ denote the minimal resolutions of E_X and G_X .*

Proof. Since G_X has Du Val singularities, then it is canonical and hence (X, G_X) is canonical. But then (X, E_X) is canonical cf. [25, §4] and so E_X is Du Val. The remaining assertion follows easily. □

2.4. Divisorial extractions.

Theorem 2.9. *Let $P \in X$ be a point of index $r > 1$.*

- (1) *There is a weighted blow up $Y \rightarrow X$ at $P \in X$ with discrepancy $1/r$.*
- (2) *Extremal divisorial contractions to $P \in X$ with discrepancy $1/r$ are given by the weighted blow-ups classified in [8], [9].*
- (3) *Extremal divisorial contractions to $P \in X$ with discrepancy greater than $1/r$ are classified in [21].*

(4) *There is a partial resolution*

$$X_n \rightarrow X_{n-1} \rightarrow \dots \rightarrow X_0 = X,$$

such that X_n has Gorenstein singularities and each map is a weighted blowup over a singular point of index $r_i > 1$ as in (1).

We have that $n \leq \rho(\tilde{E}_X/E_X)$ where $E_X \in |-K_X|$ is general and $\tilde{E}_X \rightarrow E_X$ is the minimal resolution.

Proof. See [14], [8] and [9]. $\square$

Definition 2.10. In the sequel, a *w-morphism* will denote an extremal divisorial contraction to a point $P \in X$ of index $r > 1$ with discrepancy $1/r$ see (2.9). A partial resolution as in (4) of (2.9) will be called a *w-resolution* of $P \in X$. We define $\text{dep}(X, P)$ the *depth* of X at a point $P \in X$ to be the minimum length of any *w-resolution* of (a neighborhood) of $P \in X$.

Remark 2.11. We have $\text{dep}(X, P) = 0$ if and only if X is Gorenstein at P . We have that the sum of $\text{dep}(X, P)$ for $P \in X$ is bounded below by the difficulty of X (an invariant used to show termination of flips) however, the inequality can be strict cf. [10, 2.11].

Lemma 2.12. *If $P \in X$ is a cyclic point of index r , then $\text{dep}(X) = r - 1$.*

Proof. The only *w-morphisms* are given by weighted blow ups $f : Y \rightarrow X$ such that Y contains (at most) two quotient singularities of indices a and $r - a$. The proof now follows easily by induction on the index. $\square$

Proposition 2.13. *If $(P \in X)$ is of cA/r type, then $\Xi(P \in X) - aw(P \in X) \leq \text{dep}(P \in X) \leq \Xi(P \in X) - 1$.*

Proof. We follow [8]. The singularity $P \in X$ is given by the equation $\varphi = xy + g(z^r, u) = 0$ with weights $\frac{1}{r}(\beta, -\beta, 1, r)$. Write $g(z^r, u) = \sum a_{i,j} z^{ri} u^j$, then we let

$$\lambda := aw(P \in X) = \min\{j | a_{0,j} \neq 0\},$$

and for any $s > 0$, we let

$$\nu_s := \min\{si + j | a_{i,j} \neq 0\}.$$

Notice that $\nu_1 \leq \nu_2 \leq \dots$ and $\nu_s = \lambda$ for any $s \geq \lambda$. Let

$$t := \min\{s > 0 | \nu_s = \lambda\},$$

then clearly $1 \leq t \leq \lambda$. We will show that

$$\text{dep}(P \in X) = \lambda r - t.$$

The only *w-morphisms* $f : X_1 \rightarrow X$ are weighted blow ups with weights $\frac{1}{r}(r_1, r_2, 1, r)$ such that $r_1 + r_2 = r\nu_1$ [8, §6]. We may assume that $f : X_1 \rightarrow X$ is the first *w-morphism* in a *w-resolution* of $P \in X$ computing $\text{dep}(X)$. One computes that on X_1 there are two quotient

singularities Q_1 of type $\frac{1}{r_1}(r, -1, -r)$ and Q_2 of type $\frac{1}{r_2}(r, -1, -r)$ and (possibly) a singularity Q_4 of type cA/r defined by

$$xy + g(z^r u, u)/u^{\nu_1} = xy + \sum a_{i,j} z^{ri} u^{i+j-\nu_1} = 0$$

with weights $\frac{1}{r}(\beta, -\beta, 1, r)$. If $\lambda = \nu_1$ then $Q_4 \notin X_1$ and $\text{dep}(P \in X) = r_1 + r_2 - 1 = \nu_1 r - 1$. Otherwise

$$\text{dep}(P \in X) = \text{dep}(X_1) + 1 = \nu_1 r - 1 + \text{dep}(Q_4 \in X).$$

Notice that

$$\begin{cases} \lambda(Q_4 \in X_1) = aw(Q_4 \in X_1) := \min\{j - \nu_1 | a_{0,j} \neq 0\} = \lambda - \nu_1 \\ \nu_{s-1}(Q_4 \in X_1) = \min\{(s-1)i + (i+j-\nu_1) | a_{i,j} \neq 0\} = \nu_s - \nu_1 \end{cases}$$

So $t(Q_4 \in X_1) = t - 1$. By induction on aw , we have $\text{dep}(Q_4 \in X_1) = (\lambda - \nu_1)r - (t - 1)$ and hence $\text{dep}(P \in X) = \lambda r - t$ the assertion now follows as $\lambda r = aw(P \in X)r = \Xi(P \in X)$. $\square$

Proposition 2.14. *We have the following*

- (1) *if $(P \in X)$ is of type $cAx/4$, then $\text{dep}(P \in X) \leq \Xi(P \in X) - 1$,*
- (2) *if $(P \in X)$ is of type $cD/2$ or $cD/3$ then $\text{dep}(P \in X) \leq \Xi(P \in X)$, and*
- (3) *if $(P \in X)$ is of type $cE/2$, then $\text{dep}(P \in X) \leq \Xi(P \in X) + 1$.*

Proof. This follows immediately from (4) of (2.9) and (2.1). $\square$

Proposition 2.15. *Let $f : Y \supset E \rightarrow X \ni P$ be (the germ of) a divisorial contraction to a point. Then*

$$\text{dep}(Y) \geq \text{dep}(X) - 1.$$

Proof. We follow the classification of [21]. There are several cases to consider.

If $P \in X$ is Gorenstein, then $\text{dep}(P \in X) = 0$ and the claim is clear.

If f is an w -morphism, then by definition of $\text{dep}(X)$, we have $\text{dep}(X) \leq \text{dep}(Y) + 1$ as required.

If $P \in X$ is of type cA/r , then f is a weighted blow up of discrepancy a/r (cf. [21, Thm. 1.2.i]). With the notation of Proposition (2.13), then this is obtained by using a weight $\frac{1}{r}(r_1, r_2, a, r)$ such that $r_1 + r_2 \equiv 0 \pmod{ar}$,

$$\frac{r_1 + r_2}{r} = \tau - wt(g) = \nu_a$$

where $\tau - wt(z, u) = (\frac{a}{n}, 1)$, and $z^{\frac{r_1+r_2}{a}} = z^{\frac{r\nu_a}{a}} \in g$. It follows that $\nu_s \leq \frac{s\nu_a}{a} < \nu_a$ for all $s < a$ and therefore $t \geq a$.

There are two quotient singularities $Q_1, Q_2 \in Y$ of index r_1 and r_2 and possibly a cA/r point $Q_4 \in Y$ defined by

$$\{xy - g(z^r u^a, u)/u^{\nu_a} = 0\}/\mathbb{Z}_r(-r_1, -r_2, -a, r)$$

where $g(z^r u^a, u)/u^{\nu_a} = \sum a_{i,j} z^{r_i} u^{a_i+j-\nu_a}$. It follows that

$$\sum \text{dep}(Q_i \in Y) = r_1 - 1 + r_2 - 1 + \text{dep}(Q_4 \in Y) = r\nu_a - 2 + \text{dep}(Q_4 \in Y).$$

Proceeding as in (2.13), one sees that $\text{dep}(Q_4 \in Y) = (\lambda - \nu_a)r - (t - a)$ and hence

$$\text{dep}(Y) = \lambda r - t + a - 2 = \text{dep}(X) + a - 2 \geq \text{dep}(X) - 1.$$

The remaining cases are classified in [21, 1.2, 1.3]. If $P \in X$ is not of the type cA/n and the discrepancy is $a/n > 1/n$, then there are 6 cases. Two cases of ordinary type with arbitrarily large discrepancies which are given by weighted blow-ups (cf. [21, Theorem 1.2.ii]), and 4 cases of exceptional type (cf. [21, Table 3]). We would like to remark that in general, there may be some *hidden non-Gorenstein point* on Y (i.e. non-Gorenstein points on Y at which the exceptional divisor is Cartier cf. [21, pg. 59]). However, such hidden non-Gorenstein points do not occur in the cases of exceptional type (e1, e2, e11) because we have $a/n = 4/2$ or $2/2$ and thus $2K_Y \sim_X 4E$ or $2K_Y \sim_X 2E$. It follows that if E is Cartier at a point $Q \in Y$ then K_Y is numerically equivalent to a Cartier divisor and so K_Y is Cartier at $Q \in Y$. Hence, Y is Gorenstein at Q . Therefore the baskets of Y and X are completely described by [21, 1.3].

Recall that by Reid's singular Riemann-Roch formula we have that

$$\chi(2K_X) = \frac{1}{2}K_X^3 - 3\chi(\mathcal{O}_X) + \sum_{\mathcal{B}(X)} \frac{b_i(r_i - b_i)}{2r_i},$$

and therefore

$$\chi(2K_Y) - \chi(2K_X) = \frac{1}{2}\left(\frac{a}{n}E\right)^3 + \sum_{\mathcal{B}(Y)} \frac{b_i(r_i - b_i)}{2r_i} - \sum_{\mathcal{B}(X)} \frac{b_i(r_i - b_i)}{2r_i}.$$

Claim 2.16. *Let $f : Y \supset E \rightarrow X \ni P$ be a divisorial contraction of terminal 3-folds such that $a(E, X) = 1$ (resp. $a(E, X) = 2$). Then*

$$\chi(\mathcal{O}_Y(2K_Y)) \geq \chi(\mathcal{O}_X(2K_X)) + 1.$$

Proof. If $K_Y = f^*K_X + E$, then $R^k f_* \mathcal{O}_Y(2K_Y - E) = 0$ for $k > 0$ by Kawamata-Viehweg vanishing. Since Y is terminal it is Cohen-Macaulay and hence so is E cf. [29, 5.25]. Therefore $K_E = (K_Y + E)|_E$ and we have a short exact sequence (cf. [29, 5.26])

$$0 \rightarrow \mathcal{O}_Y(2K_Y - E) \rightarrow \mathcal{O}_Y(2K_Y) \rightarrow \mathcal{O}_E(K_E) \rightarrow 0.$$

Since $-K_E = -2E|_E$ is ample, $h^1(\mathcal{O}_E(K_E)) = 0$ and so $R^1 f_* \mathcal{O}_Y(2K_Y) = 0$. Moreover, we have $R^2 f_* \mathcal{O}_Y(2K_Y) \cong R^2 f_* \mathcal{O}_E(K_E) \cong H^2(\mathcal{O}_E(K_E)) \cong \mathbb{C}$. By a Leray spectral sequence computation, we obtain

$$\begin{aligned} \chi(\mathcal{O}_Y(2K_Y)) &= \sum (-1)^i \chi(R^i f_* \mathcal{O}_Y(2K_Y)) = \chi(f_* \mathcal{O}_Y(2K_Y)) + \chi(R^2 f_* \mathcal{O}_Y(2K_Y)) \\ &= \chi(f_* \mathcal{O}_Y(2K_Y)) + h^0(R^2 f_* \mathcal{O}_Y(2K_Y)) = \chi(\mathcal{O}_X(2K_X)) + 1. \end{aligned}$$

If $K_Y = f^*K_X + 2E$, then by Kawamata-Viehweg vanishing, $R^k f_* \mathcal{O}_Y(2K_Y - 2E) = 0$ for $k > 0$. By Serre duality we have

$$h^1(\mathcal{O}_E(2K_Y)) = h^1(\mathcal{O}_E(4E)) = h^1(\mathcal{O}_E(K_E - 4E)).$$

By Kawamata-Viehweg vanishing, one sees that $h^1(\mathcal{O}_E(K_E - 4E)) = 0$ and hence $h^1(\mathcal{O}_E(2K_Y)) = 0$.

Similarly $h^1(\mathcal{O}_E(2K_Y - E)) = 0$. From the short exact sequences

$$0 \rightarrow \mathcal{O}_Y(2K_Y - (l+1)E) \rightarrow \mathcal{O}_Y(2K_Y - lE) \rightarrow \mathcal{O}_E(2K_Y - lE) \rightarrow 0$$

for $l \in \{1, 0\}$, one sees that $R^1 f_* \mathcal{O}_Y(2K_Y - E) = R^1 f_* \mathcal{O}_Y(2K_Y) = 0$ and $\mathbb{C} \cong R^2 f_* \mathcal{O}_Y(2K_Y - E) \hookrightarrow R^2 f_* \mathcal{O}_Y(2K_Y)$. The required inequality follows similarly to the previous case. $\square$

e1. f is of type $e1$ with $\frac{a}{n} = \frac{4}{2}$. We have that $\frac{4}{2}E^3 = \frac{4}{r}$ by Table 2 of [21] (compare with Table 3, where $r = 2r'$). Notice also that there is only a quotient singularity $Q \in Y$ of type $\frac{1}{2r'}(1, -1, r' - 4)$. Thus the basket of Y is $(r' - 4, 2r')$ and we have $\text{dep}(Y) = 2r' - 1$.

On the other hand $P \in X$ is a point of type $cD/2$. Notice that $\sigma(P \in X) = aw(P \in X)$ in this case and $\mathcal{B}(X) = \{aw(P \in X) \times (1, 2)\}$.

We now estimate $\text{dep}(X)$. By the singular Riemann-Roch formula, we can compute:

$$\chi(2K_Y) - \chi(2K_X) = \frac{1}{2}8E^3 + \frac{(r' - 4)(r' + 4)}{4r'} - \frac{1}{4}aw(P \in X).$$

Since $\chi(2K_Y) - \chi(2K_X) \geq 1$ cf. (2.16) and $E^3 = 2/r$, we have $r' - 1 \geq aw(p \in X)$. Therefore, by (2.14)

$$\text{dep}(X) \leq \Xi(X) = 2aw(P \in X) \leq 2r' - 2 = \text{dep}(Y) - 1$$

as required.

f is of type $e1$ with $\frac{a}{n} = \frac{2}{2}$. We have that $\frac{2}{2}E^3 = \frac{4}{r}$ with $r = 2r'$, $P \in X$ is a point of the type $cD/2$ and $Q \in Y$ is a quotient singularity of type $\frac{1}{2r'}(1, -1, r' - 2)$. We thus have $\text{dep}(Y) = 2r' - 1$.

By the singular Riemann-Roch formula, we have

$$\chi(2K_Y) - \chi(2K_X) = \frac{1}{2}E^3 + \frac{(r' - 2)(r' + 2)}{4r'} - \frac{1}{4}aw(P \in X).$$

Since $\chi(2K_Y) - \chi(2K_X) \geq 1$ cf. (2.16), we have that $r' - 1 \geq aw(P \in X)$ so that

$$\text{dep}(X) \leq \Xi(X) = 2aw(P \in X) \leq 2r' - 2 = \text{dep}(Y) - 1$$

as required.

e2. f is of type $e2$ with $\frac{a}{n} = \frac{2}{2}$. We have that $\frac{2}{2}E^3 = \frac{2}{r}$ with $r = 2r'$ and $P \in X$ is a point of the type $cD/2$. There is a singularity $Q \in Y$ of type $cA/2r'$ deforming to $2 \times \frac{1}{2r'}(1, -1, r' - 1)$. We thus have $\text{dep}(Y) \in \{4r' - 1, 4r' - 2\}$.

By the singular Riemann-Roch formula, we compute:

$$\chi(2K_Y) - \chi(2K_X) = \frac{1}{2}E^3 + 2\frac{(r' - 1)(r' + 1)}{4r'} - \frac{1}{4}aw(P \in X).$$

Since $\chi(2K_Y) - \chi(2K_X) \geq 1$ cf. (2.16), we thus have $2r' - 1 \geq aw(P \in X)$. Hence

$$\text{dep}(X) \leq \Xi(X) = 2aw(P \in X) \leq 4r' - 2 \leq \text{dep}(Y)$$

and we are done.

e11. If f is of type $e11$, then $P \in X$ is of type $cE/2$ and so $\text{dep}(X) \leq 7$. There are two quotient singularities $Q_1, Q_2 \in Y$ of type $\frac{1}{2}(1, 1, 1)$ and $\frac{1}{6}(1, -1, -1)$ respectively. Thus $\text{dep}(Y) = 6 \geq \text{dep}(X) - 1$ as required.

o3. The remaining case to consider is classified in [21, 1.2.ii]. f is a divisorial contraction to a point $P \in X$ of type $cD/2$ with discrepancy $a/2 > 1/2$. f is described explicitly as a weighted blow up and the required inequality follows from a direct computation which we describe below.

We first consider the case in [21, 1.2.ii.a], so we assume that the embedding is given by $\varphi = 0$ where

$$\begin{aligned} \varphi &= u^2 + y^2z + g(x^2, z) + uxq(x^2, z) + \lambda yx^{2\alpha-1} \\ &= u^2 + y^2z + \sum a_{ij}x^{2i}z^j + u \sum b_{ij}x^{2i+1}z^j + \lambda yx^{2\alpha-1} \end{aligned}$$

and $\mathbb{Z}_2$ acts with weights $\frac{1}{2}(1, 1, 0, 1)$.

The corresponding divisorial contraction is given by a weighted blowup $Y \rightarrow X$ with weights $\sigma_a = \frac{1}{2}(a, r, 2, r+2)$. We have that $a|(r+1)$, $a \equiv r \equiv 1 \pmod{2}$, and so we may assume that $r+1 = 2ad$. We may also assume that $a \geq 3$. Since the weight of φ with respect to σ_a is $r+1$, we have

$$\begin{cases} \min\{ai + j | a_{ij} \neq 0\} \geq r+1 = 2ad, \\ \min\{(2i+1)a + 2j | b_{ij} \neq 0\} \geq r = 2ad-1, \\ (2\alpha-1)a \geq r+2 = 2ad+1. \end{cases} \quad (\ddagger)$$

One sees that Q_2 (resp. Q_4) is a cyclic quotient point of index r (resp. $r+2$). Q_3 is a point of index 2 given by

$$\begin{aligned} \bar{\varphi} &= u^2z + y^2 + \bar{g}(x^2, z) + ux\bar{q}(x^2, z) + \lambda yx^{2\alpha-1}z^{\bar{\delta}} \\ &= u^2z + y^2 + \sum a_{ij}x^{2i}z^{\bar{\beta}_{ij}} + u \sum b_{ij}x^{2i+1}z^{\bar{\gamma}_{ij}} + \lambda yx^{2\alpha-1}z^{\bar{\delta}} \end{aligned} \quad (\dagger)$$

where

$$\begin{cases} \bar{\beta}_{ij} &= ai + j - r - 1, \\ \bar{\gamma}_{ij} &= \frac{2ai + a + 2j - r}{2}, \\ \bar{\delta} &= \frac{2a\alpha - a - r - 2}{2}. \end{cases}$$

Claim. Q_3 is a terminal singularity of type $cD/2$.

To see this, we use the coordinates $\bar{u} := y + \frac{\lambda}{2}x^{2\alpha-1}z^{\bar{\delta}}$ and $\bar{y} := u + \frac{1}{2}\sum_{\bar{\gamma}_{ij} \neq 0} b_{ij}x^{2i+1}z^{\bar{\gamma}_{ij}-1}$. Then we have

$$\bar{\varphi} = \bar{u}^2 + \bar{y}^2z + \bar{y}\left(\sum_{\bar{\gamma}_{ij}=0} b_{ij}x^{2i+1}\right) + \bar{g}(x^2, z).$$

for some $\bar{g}$. We may assume that $(\sum_{\bar{\gamma}_{ij}=0} b_{ij}x^{2i+1}) = \bar{\lambda}\bar{x}^\epsilon$ for some $\epsilon > 0$ and $\bar{\lambda} \in \mathbb{C}$ by replacing x by a suitable coordinates $\bar{x}$. Thus, this is a terminal singularity of type $cD/2$. $\square$

Notice that Q_1 is possibly a singular point of index a . However, since all hidden non-Gorenstein points are of index 2, and $a \geq 3$ is an odd integer, Q_1 is not a singular point of Y . Therefore, there is a term of the form x^b in φ with $wt(x^b) = r + 1$. In other words, $x^{4d} \in \varphi$. In conclusion, we have that

$$\text{dep}(Y) = (r - 1) + (r + 1) + \text{dep}(Q_3).$$

We will now compute $\text{dep}(X)$ by comparing with $\text{dep}(Y)$. We will construct a sequence of weighted blow ups

$$Z_a \rightarrow \dots \rightarrow Z_{k+1} \rightarrow Z_k \rightarrow \dots \rightarrow Z_0 = X$$

such that

- (1) each map $\phi_k : Z_{k+1} \rightarrow Z_k$ is a weighted blowup with weights $\sigma = \sigma_{\text{ev}} := \frac{1}{2}(1, 2d - 1, 2, 2d + 1)$ if k is even and $\sigma = \sigma_{\text{odd}} := \frac{1}{2}(1, 2d + 1, 2, 2d - 1)$ if k is odd;
- (2) moreover, let U_k^i be the standard affine piece of the weighted blowup $Z_k \rightarrow Z_{k-1}$ and R_k^i be the origin of U_k^i . The weighted blowup $\phi_k : Z_{k+1} \rightarrow Z_k$ center at R_k^3 .

We need the following claims.

Claim 1. On Z_k , there are two cyclic singularities R_k^+, R_k^- of index $2d + 1, 2d - 1$ respectively such that $\{R_k^+, R_k^-\} = \{R_k^2, R_k^4\}$.

Claim 2. On Z_k , there is a $cD/2$ singularity R_k^3 and $R_a^3 \in U_a^3 \cong U_3 \ni Q_3$.

Claim 3. Z_k is terminal and $Z_k - \{R_k^2, R_k^3, R_k^4\}$ is Gorenstein for all k .

Claim 4. $\phi_k : Z_{k+1} \rightarrow Z_k$ is a w -morphism for all k .

Grant these claims for the time being. We then have that

$$\begin{aligned} \text{dep}(X) &\leq a + \text{dep}(R_a^3) + \sum_{k=1}^a (\text{dep}(R_k^+) + \text{dep}(R_k^-)) \\ &= a + \text{dep}(Q_3) + a(2d - 2 + 2d) = a + \text{dep}(Q_3) + 4ad - 2a \\ &= \text{dep}(Q_3) + 2r + 2 - a = \text{dep}(Y) + 2 - a. \end{aligned}$$

Hence $\text{dep}(Y) \geq \text{dep}(X) + a - 2 \geq \text{dep}(X) + 1$ as required.

Proof of Claims 1, 2, 3 and 4. One can check that the equation φ_k of $Z_k \cap U_k^3$ has $\sigma - wt(\varphi_k) = 2d$ and can be written as

$$\begin{aligned} u^2 z + y^2 + \sum a_{ij} x^{2i} z^{\beta_{ij}(k)} + u \sum b_{ij} x^{2i+1} z^{\gamma_{ij}(k)} + \lambda y x^{2\alpha-1} z^{\delta(k)}, & \text{ if } k \text{ is odd} \\ u^2 + y^2 z + \sum a_{ij} x^{2i} z^{\beta_{ij}(k)} + u \sum b_{ij} x^{2i+1} z^{\gamma_{ij}(k)} + \lambda y x^{2\alpha-1} z^{\delta(k)}, & \text{ if } k \text{ is even.} \end{aligned}$$

Where $\beta_{ij}(0) = j$ and $\beta_{ij}(k + 1) = \beta_{ij}(k) + i - 2d$ so that

$$\beta_{ij}(k) = k(i - 2d) + j;$$

$\gamma_{ij}(0) = j$ and $\gamma_{ij}(k) = \gamma_{ij}(k-1) + \frac{2d+1}{2} + \frac{2i+1}{2} - 2d$ if k is odd and $\gamma_{ij}(k) = \gamma_{ij}(k-1) + \frac{2d-1}{2} + \frac{2i+1}{2} - 2d$ if k is even, so that

$$\begin{aligned}\gamma_{ij}(k) &= k\frac{2i+1}{2} - kd + j, & \text{if } k \text{ is even} \\ \gamma_{ij}(k) &= k\frac{2i+1}{2} - kd + j + \frac{1}{2}, & \text{if } k \text{ is odd;} \end{aligned}$$

$\delta(0) = 0$ and $\delta(k) = \delta(k-1) + \frac{2d-1}{2} + \frac{2\alpha-1}{2} - 2d$ if k is odd and $\delta(k) = \delta(k-1) + \frac{2d+1}{2} + \frac{2\alpha-1}{2} - 2d$ if k is even, so that

$$\begin{aligned}\delta(k) &= k\frac{2\alpha-1}{2} - kd, & \text{if } k \text{ is even} \\ \delta(k) &= k\frac{2\alpha-1}{2} - kd - \frac{1}{2}, & \text{if } k \text{ is odd.} \end{aligned}$$

We need to verify that $\sigma - wt(\varphi_k) = 2d$ with respect to the weight $\sigma_{\text{odd}} = \frac{1}{2}(1, 2d+1, 2, 2d-1)$ or $\sigma_{\text{ev}} = \frac{1}{2}(1, 2d-1, 2, 2d+1)$ (depending on the parity of k). One can easily see that $\sigma_{\text{odd}} - wt(u^2z) = 2d$ and $\sigma_{\text{odd}} - wt(y^2) = 2d+1$ (resp. $\sigma_{\text{ev}} - wt(u^2) = 2d+1$ and $\sigma_{\text{ev}} - wt(y^2z) = 2d$) if k is odd (resp. even). Moreover, since $x^{4d} \in \varphi$, we have $x^{4d} \in \varphi_k$ for all k . Therefore, $wt(\varphi_k) \leq 2d$. One sees that if $\beta_{ij}(k+1) \geq 0$, $\gamma_{ij}(k+1) \geq 0$ and $\delta(k+1) \geq 0$, then $wt(\varphi_k) \geq 2d$.

Therefore it is enough to show that $\beta_{ij}(k) \geq 0$, $\gamma_{ij}(k) \geq 0$ and $\delta(k) \geq 0$ for all $1 \leq k \leq a$ and all i and j with $a_{i,j} \neq 0$, resp. $b_{i,j} \neq 0$.

Recall that by $(\ddagger)$, $ai + j \geq 2ad$, thus

$$\beta_{ij}(k) = k(i-2d) + j = \frac{k}{a}(ai+j) - \frac{kj}{a} - 2kd + j \geq j - \frac{kj}{a} = j\frac{a-k}{a} \geq 0.$$

Next, recall that by $(\ddagger)$, $(2i+1)a + 2j \geq 2ad - 1$, hence

$$\begin{aligned}\gamma_{ij}(k) &\geq k\frac{2i+1}{2} - kd + j = \frac{k}{2a}(2i+1)a - kd + j \\ &\geq \frac{k}{2a}(2ad - 1 - 2j) - kd + j = \frac{2aj - 2kj - k}{2a} \geq \frac{-k}{2a} \geq \frac{-1}{2}.\end{aligned}$$

Also, recall that by $(\ddagger)$, $(2\alpha-1)a \geq 2ad + 1$, hence

$$\begin{aligned}\delta(k) &\geq k\frac{2\alpha-1}{2} - kd - \frac{1}{2} = \frac{k}{2a}(2\alpha-1)a - kd - \frac{1}{2} \\ &\geq \frac{k}{2a}(2ad + 1) - kd - \frac{1}{2} = \frac{k}{2a} - \frac{1}{2} \geq \frac{-1}{2}.\end{aligned}$$

The above inequalities now follow because $\beta_{ij}(k)$ and $\delta(k)$ are integers.

Note that $\beta_{ij}(a) = \bar{\beta}_{ij}$, $\gamma_{ij}(a) = \bar{\gamma}_{ij}$ and $\delta(a) = \bar{\delta}$. Thus $R_a^3 \in Z_a$ is indeed isomorphic to $Q_3 \in Y$.

Claims 1 and 2 now follow easily from the equations φ_k .

Claim 3 also follows by explicit computation.

We now verify that the homogeneous part $\varphi_{k, \sigma - wt = 2d}$, which defines the exceptional divisor E_k in $\mathbb{P}(1, 2d+1, 2, 2d-1)$ (resp. $\mathbb{P}(1, 2d-1, 2, 2d+1)$), is irreducible. To see this, notice that as we have seen $x^{4d} \in \varphi$, and so $a_{2d,0} \neq 0$. Since $\beta_{2d,0}(k) = 0$, one sees that $x^{4d} \in \varphi_k$ for all k . If k is odd, we see that $u^2z, x^{4d} \in \varphi_{k, \sigma_{\text{odd}} - wt = 2d}$. Hence $\varphi_{k, \sigma_{\text{odd}} - wt = 2d}$

is irreducible. If k is even, we see that $y^2z, x^{4d} \in \varphi_{k, \sigma_{\text{ev}} - wt = 2d}$. Hence $\varphi_{k, \sigma_{\text{ev}} - wt = 2d}$ is irreducible. Since $wt(\varphi) = 2d$, we thus have

$$K_{Z_{k+1}} = \phi_k^* K_{Z_k} + (wt(xyzu) - wt(\varphi_k) - 1)E_{k+1} = \phi_k^* K_{Z_k} + \frac{1}{2}E_{k+1}.$$

This completes the proof. $\square$

The other case with discrepancy $\frac{a}{2}$ can be treated similarly. We may assume that $r + 2 = (2d + 1)a$ for some $d > 0$. The local equation of $P \in X$ is given by

$$\begin{cases} \phi_1 : & = u^2 + yw + p(x^2, z) = 0 \\ & = u^2 + yw + \sum a_{ij}x^{2i}z^j = 0 \\ \phi_2 : & = yz + x^{2d+1} + xzq(x^2, z) + w = 0 \\ & = yz + x^{2d+1} + \sum b_{ij}x^{2i+1}z^{j+1} + w = 0 \end{cases}$$

The divisorial contraction $Y \rightarrow X$ with discrepancy $\frac{a}{2}$ is a weighted blowup with weight $\sigma_a = \frac{1}{2}(a, r, 2, r + 2, r + 4)$ with respect to the coordinates (x, y, z, u, w) . Thus the local equation for Y near Q_3 is

$$\begin{cases} \phi_{1,a} & = u^2 + yw + \sum a_{ij}x^{2i}z^{ai+j-(r+2)} = 0 \\ \phi_{2,a} & = y + x^{2d+1} + \sum b_{ij}x^{2i+1}z^{j+1+\frac{(2i+1)a-(r+2)}{2}} + wz = 0 \end{cases}$$

Similarly to the previous case, we construct a sequence of weighted blow-ups $Z_a \rightarrow Z_{a-1} \rightarrow \dots \rightarrow Z_1 \rightarrow Z_0 = X$ by weights $\sigma_{\text{ev}} = \frac{1}{2}(1, 2d - 1, 2, 2d + 1, 2d + 3)$ and $\sigma_{\text{odd}} = \frac{1}{2}(1, 2d + 3, 2, 2d + 1, 2d - 1)$, where $Z_{k+1} \rightarrow Z_k$ is a weighted blowup with weight σ_{ev} (resp. σ_{odd}) if k is even (resp. odd).

By a direct computation, one sees that if k is odd, then the local equations are

$$\begin{cases} \phi_{1,k} & = u^2 + yw + \sum a_{ij}x^{2i}z^{ki+j-k(2d+1)} = 0 \\ \phi_{2,k} & = y + x^{2d+1} + \sum b_{ij}x^{2i+1}z^{j+1+\frac{k(2i+1)-k(2d+1)}{2}} + wz = 0 \end{cases}$$

If k is even, then the local equations are

$$\begin{cases} \phi_{1,k} & = u^2 + yw + \sum a_{ij}x^{2i}z^{ki+j-k(2d+1)} = 0 \\ \phi_{2,k} & = yz + x^{2d+1} + \sum b_{ij}x^{2i+1}z^{j+1+\frac{k(2i+1)-k(2d+1)}{2}} + w = 0 \end{cases}$$

Notice that we have $wt(\phi_1) = 2d + 1$ (resp. $wt(\phi_2) = \frac{2d+1}{2}$) because $ki + j - k(2d + 1) \geq 0$ (resp. $j + 1 + \frac{k(2i+1)-k(2d+1)}{2} \geq 0$) for all $1 \leq k \leq a$ and all i and j such that $a_{i,j} \neq 0$ (resp. $b_{i,j} \neq 0$).

We now compare the singularities. On Y , there are cyclic quotient points Q_2 and Q_5 of index r and $r + 4$ respectively and a $cD/2$ point Q_3 . On Z_k , there are two cyclic quotient point R_k^+, R_k^- of index $2d + 3, 2d - 1$ respectively and a $cD/2$ point R_k^3 . The point $R_a^3 \in Z_a$ has index 2 and is isomorphic to Q_3 in Y . Hence we conclude that

$$\text{dep}(X) \leq a + \text{dep}(R_a^3) + \sum_{k=1}^a (\text{dep}(R_k^+) + \text{dep}(R_k^-))$$

$$\begin{aligned}
&= a + \text{dep}(Q_3) + a(2d + 2 + 2d - 2) = a + \text{dep}(Q_3) + 4ad \\
&= \text{dep}(Q_3) + 2r + 4 - a = \text{dep}(Y) + 2 - a.
\end{aligned}$$

Since $a \geq 3$, we have $\text{dep}(Y) \geq \text{dep}(X) + a - 2 \geq \text{dep}(X) + 1$. $\square$

3. DECOMPOSITION

The purpose of this section is to prove the main result (1.1). We begin by proving the following key result.

Theorem 3.1. *Let $g : X \supset C \rightarrow W \ni R$ be an extremal neighborhood. If X is not Gorenstein, then there exists a w -morphism $f : Y \rightarrow X$, such that $C_Y \cdot K_Y \leq 0$, where C_Y denotes the proper transform of C in Y .*

Proof. We use the classification of extremal neighborhoods. Each case is elementary.

Let $P \in C \subset X$ be an isolated terminal singularity of index $r > 1$. By (2.9), there is a weighted blowup $f : Y \rightarrow X$ with exceptional divisor F such that $K_Y = f^*K_X + \frac{1}{r}F$. We will need to compute

$$(2) \quad C_Y \cdot K_Y = C_Y \cdot (f^*K_X + \frac{1}{r}F) = C \cdot K_X + \frac{1}{r}C_Y \cdot F.$$

We follow the classification summarized in Appendix A and B of [34] together with the classification of the terminal singular point P cf. [30, pg. 541], and we will apply the results of [8] and [9].

Suppose that $C \not\subset E_X$ where E_X is a general section of $|-K_X|$. By (2.2), E_X is normal with Du Val singularities. By (2.7), we have that $f^*(K_X + E_X) = K_Y + E_Y$ where $E_Y = f_*^{-1}E_X$. Therefore

$$0 = C \cdot (K_X + E_X) = C_Y \cdot (K_Y + E_Y) \geq C_Y \cdot K_Y.$$

It suffices therefore to consider those extremal neighborhoods with $C \subset E_X$. By [30, 2.2] and [35], the only cases to consider are IC, IIB, exceptional IA + IA, IA + IA + III and semistable IA + IA.

IC. By [34, pg. 243], we have that

$$C^\# = \{y_1^{r-2} - y_2^2 = y_4 = 0\} \subset$$

$$X^\# = \{y_1, y_2, y_4\}/\mathbb{Z}_r(2, r-2, 1)$$

where $r \geq 5$ is odd. Following [8, 3.9], if we blow up $\mathbb{C}^3/\mathbb{Z}_r(2, r-2, 1)$ with weight $\frac{1}{r}(2, r-2, 1)$, we obtain a divisorial extraction with discrepancy $1/r$.

Computing on U_1 yields $C_Y = \{1 - \bar{y}_2^2 = \bar{y}_4 = 0\}$ and $F = \{\bar{y}_1 = 0\}$ where $\mathbb{Z}_2$ acts with weights $(r, -r+2, -1)$. Since r is odd, on U_1 we have $F \cap C_Y = Q$ is a point with multiplicity 1.

Computing on U_2 yields $C_Y = \{\bar{y}_1^{r-2} - 1 = \bar{y}_4 = 0\}$ and $F = \{\bar{y}_2 = 0\}$ where $\mathbb{Z}_{r-2}$ acts with weights $(-2, r, -1)$. So on U_2 we have $F \cap C_Y = Q$ where $Q \in U_1 \cap U_2$ is the same point as above.

Computing on U_3 yields no further intersection points. Therefore, $C_Y \cdot F = 1$ and hence $K_Y \cdot C_Y \leq 0$.

IIB. We have $cAx/4$ point $P \in C$. By [34, pg. 243], we have

$$C^\sharp = \{y_1^2 - y_2^3 = y_3 = y_4 = 0\} \subset$$

$$X^\sharp = \{y_1^2 - y_2^3 + g_3y_3 + g_4y_4 = 0\}$$

where $\mathbb{Z}_4$ acts on $\mathbb{C}^4$ with weights $(3, 2, 1, 1)$. By [8, §7] there is a w -morphism $f : Y \supset F \rightarrow X \ni P$ which is a weighted blowup with equivalent weights $\frac{1}{4}(r_1, r_2, r_3, r_4)$ such that $r_1 \equiv 3, r_2 \equiv 2, r_3 \equiv r_4 \equiv 1 \pmod{4}$.

We distinguish the following cases:

If $2r_1 > 3r_2$, then computation on U_1 yields that $C_Y = \{\bar{y}_1^{\frac{2r_1-3r_2}{4}} - \bar{y}_2^3 = \bar{y}_3 = \bar{y}_4 = 0\}$, F is defined by $\bar{y}_1 = 0$ where $\mathbb{Z}_{r_1}$ acts with weights $(4, -r_2, -r_3, -r_4)$. Moreover, computation on U_2, U_3 and U_4 yields no further points in $C_Y \cap F$. Thus $C_Y \cdot F = \frac{3}{r_1} \leq 1$.

If $2r_1 < 3r_2$, then similar computation yields that $C_Y \cdot F = \frac{2}{r_2} \leq 1$.

Finally, if $2r_1 = 3r_2$, then a similar computation that $C_Y \cdot F = \frac{3}{r_1} = \frac{2}{r_2}$.

In conclusion we have that

$$C_Y \cdot F = \min\left\{\frac{3}{r_1}, \frac{2}{r_2}\right\} \leq 1,$$

and so $C_Y \cdot K_Y \leq 0$.

Before we turn our attention to the remaining three cases, it is necessary to prove the following.

Lemma 3.2 (Blowing up cA/r points on an extremal neighborhood of type IA). *Let $P \in C \subset X$ be a point of type cA/r on an extremal neighborhood of type IA , then by [34, pg. 243], we have*

$$C^\sharp = \{y_1^{a_2} - y_2^{a_1} = y_3 = y_4 = 0\} \subset$$

$$X^\sharp = \{\varphi = g_2(y_1^{a_2} - y_2^{a_1}) + g_3y_3 + g_4y_4 = 0\}$$

where $\mathbb{Z}_r$ acts on $\mathbb{C}^4$ with weights $(a_1, a_2, -a_1, 0)$ and $(a_1a_2, r) = 1$.

Then there is a w -morphism $Y \supset F \rightarrow X \ni P$ of discrepancy $1/r$ given by a weight $\sigma = \frac{1}{r}(r_1, 1, r_2, r)$ such that $C_Y \cdot F = a_1/r_1$ where C_Y is the strict transform of C and $r_1 \equiv -r_2 \equiv a_1a_2^{-1}$ modulo r .

Proof. Let $r > \lambda > 0$ be an integer such that $a_2\lambda = qr + 1$ for some integer $q \geq 0$ and α be the residue of $a_1\lambda$ modulo r . By [8, Lemma 6.5], there is a w -morphism $f : Y \supset F \rightarrow X \ni P$ of discrepancy $1/r$ given by a weight $\sigma = \frac{1}{r}(r_1, 1, r_2, r)$ where r_1 and r_2 are positive integers such that $r_1 \equiv -r_2 \equiv \alpha$ modulo r .

The weighted blow up is covered by four affine pieces $U_1, \dots, U_4$ whose equations are described in [8, §6]. An easy computation shows that

$F \cap C_Y = \emptyset$ on U_3 and U_4 . Since $r_1 \equiv a_1\lambda \pmod{r}$, we may write $r_1 = a_1\lambda + tr$ for some t . Hence

$$(3) \quad a_2r_1 = a_1a_2\lambda + a_2tr = (a_1q + a_2t)r + a_1.$$

Note that as $a_2r_1 > 0$ and $r > a_1 > 0$, we have that $a_1q + a_2t \geq 0$. Computation in U_2 now yields that $C_Y = \{\bar{y}_1^{a_2}\bar{y}_2^{a_1q+a_2t} - 1 = \bar{y}_3 = \bar{y}_4 = 0\}$, $F = \{\bar{y}_2 = 0\}$. Therefore $C_Y \cap F = \emptyset$ in U_2 .

Computation in U_1 yields that $C_Y = \{\bar{y}_1^{a_1q+a_2t} - \bar{y}_2^{a_1} = \bar{y}_3 = \bar{y}_4 = 0\}$, $F = \{\bar{y}_1 = 0\}$ where $\mathbb{Z}_{r_1}$ acts with weights $(r, -1, -r_2, -r)$. Therefore $C_Y \cdot F = a_1/r_1$. $\square$

Exceptional IA + IA. If g is a flipping contraction (resp. a divisorial contraction), then we have two IA points P and Q such that Q is a $cA/2$ (resp. $cA/2$ or $cAx/2$ or $cD/2$) point and P is a cA/r point of odd index $r \geq 5$ (resp. $r \geq 3$). Since (cf. §2.2)

$$C \cdot K_X = -1 + \frac{1}{2} + \frac{r - a_2}{r} = \frac{1}{2} - \frac{a_2}{r} < 0$$

we have $a_2 > \frac{r}{2}$. Let $s = 2a_2 - r > 0$. Then $C \cdot K_X = -s/2r$.

By [30, 2.13.1], we have $a_1 = 1$. We consider the w -morphism $f : Y \supset F \rightarrow X \ni P$ over the point P given by (3.2). We thus have $C_Y \cdot F = \frac{1}{r_1}$ and hence

$$K_Y \cdot C_Y = \frac{-s}{2r} + \frac{1}{rr_1} = \frac{1}{r} \left(-\frac{s}{2} + \frac{1}{r_1} \right).$$

Notice that $a_2r_1 \equiv a_1 = 1$ modulo r and so $a_2r_1 = 1 + qr$ for some integer $q \geq 0$ and hence $sr_1 = 2 + (2q - r_1)r$. Since $sr_1 > 0$ and $r > 1$, it follows that $sr_1 \geq 2$ and hence $1/r_1 - s/2 \leq 0$. Therefore, we conclude that $C_Y \cdot K_Y \leq 0$.

Semistable IA + IA. Let C be the flipping curve. Then C contains two IA points P and P' corresponding to singularities of type cA/r and cA/r' respectively. We assume that $r \geq r'$. By [30, §13, pg. 679], the curve $C^\sharp$ is given by the x -axis, and $X^\sharp$ is given by $xy - z^{dr} + tf(x, y, z, t) = 0 \subset \mathbb{C}^4$ with weights $\frac{1}{r}(1, -1, a, 0)$. It follows that $a_1 = 1$ and $a_2 = a$. Similarly for the second singular point we have $a'_1 = 1$ and $a'_2 = a'$.

Let $\frac{a}{r} + \frac{a'}{r'} = 1 + \frac{\delta}{rr'}$, then $\delta > 0$ and $K_X \cdot C = -\frac{\delta}{rr'}$. We write

$$(4) \quad ar' + a'r = rr' + \delta.$$

Let $f : Y \supset F \rightarrow X \ni P$ be the weighted blow up of X at P with weight $\sigma = \frac{1}{r}(r_1, 1, r_2, r)$ as described in (3.2). In particular $C_Y \cdot F = 1/r_1$. Recall that since $r_1 \equiv a^{-1} \pmod{r}$, we have that

$$(5) \quad ar_1 = \gamma r + 1 \text{ for some } \gamma \geq 0.$$

Suppose that $\gamma = 0$, then $ar_1 = 1$. Hence $a = r_1 = 1$. Then

$$\frac{r' - 1}{r'} \geq \frac{a'}{r'} = 1 + \frac{\delta}{rr'} - \frac{1}{r} > \frac{r - 1}{r},$$

so that $r' > r$ which is a contradiction.

Therefore, we have $\gamma > 0$. By (5) and (4), we have

$$ar_1\delta \equiv \delta \equiv ar' \pmod{r}.$$

Hence $r_1\delta \equiv r' \pmod{r}$. Since $r' \leq r$, we thus have $r_1\delta \geq r'$. It follows that

$$C_Y \cdot K_Y = -\frac{\delta}{rr'} + \frac{1}{rr_1} \leq -\frac{\delta}{rr'} + \frac{\delta}{rr'} = 0.$$

IA + IA + III. In this case $C \subset X$ is divisorial and the two IA points are both ordinary of cA type of index 2 and $r \geq 3$. By [30, 2.12.1], we also have that $a_1 = 1$ at the cA/r point. The same computation as in the Exceptional IA+IA case shows that $C_Y \cdot K_Y \leq 0$. $\square$

Theorem 3.3. *Let $g : X \supset C \rightarrow W \ni P$ be an extremal neighborhood which is isolated (resp. divisorial). If X is not Gorenstein, then we have a diagram*

$$\begin{array}{ccc} Y & \dashrightarrow & Y' \\ f \downarrow & & \downarrow f' \\ X & & X' \\ & g \searrow & \swarrow g' \\ & & W \end{array}$$

where $Y \dashrightarrow Y'$ consists of flips and flops over W , f is a w -morphism, f' is a divisorial contraction (resp. a divisorial contraction to a curve) and $g' : X' \cong X^+ \rightarrow W$ is the flip of g (resp. g' is divisorial contraction to a point).

Proof. 1. By [32, Lemma 3.15], we have that any g -exceptional curve C' is not numerically effective. In particular, there is an effective irreducible divisor $D \in \text{Div}(X)$ such that $D \cdot C' < 0$.

In the case of a divisorial contraction, this divisor D is nothing but the exceptional divisor to be contracted. In the case of small contraction, the divisor D moves in a family.

2. Let $f : Y \rightarrow X$ be the w -morphism with exceptional divisor F given by (3.1). Let l be a curve on F . Let C_Y and D_Y be the proper transforms of C and D respectively.

Since $\rho(Y/W) = 2$ and $\text{Exc}(g \circ f) = F \cup C_Y$,

$$\text{NE}(Y/W) = \mathbb{R}_+ l + \mathbb{R}_+ C_Y.$$

3. One can flip or flop along C_Y .

By 2, C_Y is extremal. Moreover, $f^*D = D_Y + \lambda F$ where $\lambda > 0$ and hence

$$C_Y \cdot (K_Y + \epsilon D_Y) \leq C_Y \cdot \epsilon D_Y = \epsilon C_Y \cdot (f^*D - \lambda F) < 0.$$

Thus, we have a $(K_Y + \epsilon D_Y)$ -flip $Y \dashrightarrow Y^+$, which is a K_Y -flip or K_Y -flop.

4. Notice that K_{Y^+} is not nef over W .

More precisely, let $l \subset F$ be a general curve, then $l \cap C_Y = \emptyset$. Hence the proper transform l_+ of l in Y^+ satisfies $l_+ \cdot K_{Y^+} = l \cdot K_Y < 0$. Moreover, the family of such curves dominates the divisor F_{Y^+} .

We now run the K_{Y^+} minimal model program over W . By termination of flips, we must end up with a divisorial contraction $Y' \rightarrow X'$. Therefore, we have the required diagram.

5. Suppose that $g : X \rightarrow W$ is divisorial with exceptional divisor D . Notice that

$$\text{Exc}(Y'/W) \supset D_{Y'} \cup F_{Y'}$$

where $D_{Y'}$ and $F_{Y'}$ are the strict transforms of F and D on Y .

We claim that f' contracts the divisor $D_{Y'}$ to a curve.

Suppose to the contrary that $Y' \rightarrow X'$ contracts $F_{Y'}$. Then we see that $X' \rightarrow W$ has relative Picard number $\rho(X'/W) = 1$ containing an exceptional divisor $D_{X'}$. Notice furthermore that D and $D_{X'}$ are isomorphic away from finitely many curves. Therefore, we have $X' \cong X$ by an easy argument cf. [18, Lemma 3.4]. By the same argument, we also have $Y \cong Y'$, which is a contradiction.

Hence $f' : Y' \rightarrow X'$ contracts $D_{Y'}$. Notice that $g(D)$ is a curve in W . Hence $g'(f'(D_{Y'}))$ is a curve. In particular, $f'(D_{Y'})$ is a curve.

Since $g(f(F))$ is a point. It follows that $g'(F_{X'})$ is a point.

6. Suppose that $g : X \rightarrow W$ is isolated. We claim that $X' \cong X^+$ is the flip.

First of all, X and X' are isomorphic in codimension 1 and the relative Picard number $\rho(X'/W)$ is equal to 1.

If $K_{X'}$ is g' -ample over W , we are done.

If $K_{X'}$ is g' -trivial, we have that $K_{X'} = g'^*K_W$. This implies that K_W is $\mathbb{Q}$ -Cartier, which is impossible.

If $K_{X'}$ is g' -negative, we see that

$$X \cong \text{Proj}_W \left(\bigoplus_{m \geq 0} \mathcal{O}_W(-mK_W) \right) \cong X'.$$

It is clear that F_Y and $F_{Y'}$ correspond to the same valuation in the function field. We thus have $Y \cong Y'$ by [18, Lemma 3.4], which is absurd because $Y \dashrightarrow Y'$ is a composition of a sequence of $(K_Y + \epsilon D_Y)$ -flips. $\square$

Remark 3.4. We would like to remark that if $g : X \rightarrow W$ is isolated, then either $f' : Y' \rightarrow X'$ is a divisorial contraction to a curve or at least one of $Y_i \dashrightarrow Y_{i+1}$ is a flip. To see this, pick any flipped curve $C_{X'}$ such that $K_{X'} \cdot C_{X'} > 0$. Suppose that f' is a divisorial contraction to a point, then $C_{X'} \not\subset f'(F_{Y'})$. Hence

$$K_{Y'} \cdot C_{Y'} = (f'^*K_{X'} + aF_{Y'}) \cdot C_{Y'} = K_{X'} \cdot C_{X'} + aC_{Y'} \cdot F_{Y'} > 0,$$

where $C_{Y'}$ denotes the proper transform of $C_{X'}$ in Y' and $a > 0$. If there are no flips $Y_i \dashrightarrow Y_{i+1}$, then $Y \dashrightarrow Y'$ is a flop and if C_Y is the proper transform of $C_{Y'}$ in Y , then $K_Y \cdot C_Y = K_{Y'} \cdot C_{Y'} > 0$. Therefore, as $f : Y \rightarrow X$ is a divisorial contraction, C_Y maps to a curve $C_X \subset X$ which is the flipping curve. But then C_Y is (by construction) the flopping curve of $Y \dashrightarrow Y'$ and so $C_Y \cdot K_Y = 0$. This is the required contradiction.

Remark 3.5. The result of (3.3) also holds if we assume that $g : X \supset C \rightarrow W \ni P$ is a flipping contraction or a divisorial contraction to a curve. In fact, by (2.3) and (2.4), g factors through an extremal neighborhood $g' : X \supset C' \rightarrow W' \ni P'$. Therefore, (3.1) also applies in this context so that we have a w -morphism $f : Y \rightarrow X$ such that $K_Y \cdot C_Y \leq 0$. The rest of the argument is the same as the proof of (2.3).

Remark 3.6. In order to prove (2.2), we will need an invariant which is minimal for Gorenstein varieties and that improves under w -morphisms and flips. There seem to be two possible approaches, the first one using general elephants $E_X \in |-K_X|$ (i.e. assuming that there exists an element $E_X \in |-K_X|$ with Du Val singularities) and the second one using $\text{dep}(X)$.

It is expected that if $g : X \rightarrow W$ is the germ of a flipping or divisorial contraction, then the general elements $E_X \in |-K_X|$ and $E_W \in |-K_W|$ have Du Val singularities (this is known in the divisorial to point case by [21], in the extremal neighborhood case by (2.2) and is expected to hold unconditionally). If we assume that E_X is Du Val, then there is a simpler proof for (1.1) not involving the invariants $\text{dep}(X)$ which we now explain.

Proof of (1.1) assuming that E_X is Du Val. We may work locally over W and so we may assume that there is a point $Q \in W$ such that $W - Q$ is smooth and $X - f^{-1}(Q)$ is smooth. Let E_X be as above and let $\tilde{E}_X$ be the minimal resolution of E_X then $\rho(\tilde{E}_X/E_X)$ defines an invariant of (the germ of) $g : X \rightarrow W$. We will proceed by induction on $\rho(\tilde{E}_X/E_X)$.

If $\rho(\tilde{E}_X/E_X) = 0$, then E_X is smooth and hence terminal. By inversion of adjunction (X, E_X) is terminal. By classification of terminal 3-fold singularities, X is Gorenstein. By (2.6), g is divisorial and it is the blow up of a LCI curve in W .

We may therefore assume that $\rho(\tilde{E}_X/E_X) > 0$. By (3.5), we may factor $X \rightarrow W$ by rational maps over W

$$X \xleftarrow{f} Y \dashrightarrow Y' \xrightarrow{f'} X'.$$

Let $f : F \subset Y \rightarrow P \in X$ be the f -exceptional divisor. By (2.7), $f^*(K_X + E_X) = K_Y + E_Y$ where $E_Y = f_*^{-1}E_X$. By (2.7), $\tilde{E}_X \rightarrow E_X$ factors through E_Y . It follows by (2.8) that $\rho(\tilde{E}_Y/E_Y) < \rho(\tilde{E}_X/E_X)$.

The rational map $Y \dashrightarrow Y'$ is given by a sequence of flips and flops $Y_i \dashrightarrow Y_{i+1}$.

Claim 3.7. *If $Y_i \dashrightarrow Y_{i+1}$ is a flip (resp. a flop), then $\rho(\tilde{E}_{Y_i}/E_{Y_i}) \geq \rho(\tilde{E}_{Y_{i+1}}/E_{Y_{i+1}})$ where E_{Y_i} denotes the strict transform of E_X and $\tilde{E}_{Y_i}$ denotes its minimal resolution.*

Proof of the Claim. By (2.3), we may assume that the flip $Y_i \dashrightarrow Y_{i+1}$ factors in the analytic category, through a sequence of flips with irreducible flipping curve. Let $\bar{Y}_i \rightarrow \bar{W}_i \leftarrow \bar{Y}_i^+$ be one of these flips. It suffices to show that $\rho(\tilde{E}_{\bar{Y}_i}/E_{\bar{Y}_i}) \geq \rho(\tilde{E}_{\bar{Y}_i^+}/E_{\bar{Y}_i^+})$. Since $\bar{Y}_i \dashrightarrow \bar{Y}_i^+$ is an isomorphism on the complement of the flipping curve $\bar{C}_i$, we may assume that $\bar{Y}_i \supset \bar{C}_i \rightarrow \bar{W}_i \ni \bar{w}_i$ is an extremal neighborhood. Since $\bar{Y}_i \dashrightarrow \bar{Y}_i^+$ is a $(K_{\bar{Y}_i} + E_{\bar{Y}_i})$ -flop, we have that $E_{\bar{Y}_i^+} \rightarrow E_{\bar{W}_i}$ is crepant i.e. $K_{E_{\bar{Y}_i^+}/E_{\bar{W}_i}} = 0$. Therefore, if $E_{\bar{Y}_i}$ has Du Val singularities, then $E_{\bar{Y}_i^+} \in |-K_{\bar{Y}_i^+}|$ is also Du Val. Moreover, the rational map $\tilde{E}_{\bar{Y}_i} \dashrightarrow E_{\bar{Y}_i^+}$ is a morphism. Since the flipping curve is irreducible, $\rho(E_{\bar{Y}_i}/E_{\bar{W}_i}) \leq 1$. Since $E_{\bar{Y}_i^+} \in |-K_{\bar{Y}_i^+}|$ and $K_{\bar{Y}_i^+}$ is ample over $\bar{W}_i$, $E_{\bar{Y}_i^+}$ contains the flipped curves so that $\rho(E_{\bar{Y}_i^+}/E_{\bar{W}_i}) \geq 1$. Therefore, we have

$$\rho(\tilde{E}_{\bar{Y}_i^+}/E_{\bar{Y}_i^+}) \leq \rho(\tilde{E}_{\bar{Y}_i}/E_{\bar{Y}_i^+}) \leq \rho(\tilde{E}_{\bar{Y}_i}/E_{\bar{Y}_i}).$$

The case of a flop is similar. In this case $K_{\bar{Y}_i} \equiv_{\bar{W}_i} 0$ and $K_{\bar{Y}_i^+} \equiv_{\bar{W}_i} 0$. If $E_{\bar{Y}_i}$ does not intersect the flopping locus then $E_{\bar{Y}_i^+}$ does not intersect the flopping locus. If $E_{\bar{Y}_i}$ intersects the flopping locus, then $E_{\bar{Y}_i^+}$ intersects the flopped locus and hence it contains the flopped locus. Therefore $\rho(\tilde{E}_{\bar{Y}_i}/E_{\bar{Y}_i^+}) \leq \rho(\tilde{E}_{\bar{Y}_i}/E_{\bar{Y}_i})$ and we conclude as in the previous case. $\square$

It follows that if $G_{Y_i} \in |-K_{Y_i}|$ is general, then it has Du Val singularities and

$$\rho(\tilde{E}_X/E_X) > \rho(\tilde{E}_{Y_i}/E_{Y_i}) \geq \rho(\tilde{G}_{Y_i}/G_{Y_i}) \quad \text{for all } i.$$

By induction, we obtain the required factorization of $Y \dashrightarrow Y'$ and of $Y' \rightarrow X'$. $\square$

We will now prove (1.1) in full generality using $\text{dep}(X)$. We will need the following propositions that we believe are of independent interest.

Proposition 3.8. *Let $X \rightarrow W$ be a flipping contraction and let $X \dashrightarrow X'$ be the flip. Then $\text{dep}(X) > \text{dep}(X')$.*

Proposition 3.9. *Let $X \rightarrow W$ be a extremal contraction to a curve. Then $\text{dep}(X) \geq \text{dep}(W)$.*

Proof of (1.1) and (3.8), (3.9). We may work locally over the base W and so we may assume that $g : X \rightarrow W$ is the germ of a flipping (resp.

a divisorial to curve) contraction. By (3.5), we may factor $X \dashrightarrow X'$ (resp. $X \rightarrow W$) by rational maps over W

$$X \xleftarrow{f} Y \dashrightarrow Y' \xrightarrow{f'} X'.$$

We proceed by induction on $\text{dep}(X)$. If $\text{dep}(X) = 0$, then X is Gorenstein. There is no flipping contraction. Therefore g is a divisorial contraction to a curve, given by blowing up a LCI curve in W . Hence the above two Propositions and Theorem 1.1 holds when $\text{dep}(X) = 0$.

We may therefore assume that $\text{dep}(X) > 0$ and the statement holds for all threefolds with $\text{dep} < \text{dep}(X)$.

By the proof of (3.1), we may assume that $Y \rightarrow X$ is the first weighted blowup of a w -resolution of X , so that $\text{dep}(X) - 1 = \text{dep}(Y)$. If $Y \dashrightarrow Y^+$ is a flop, then by [23] Y and Y^+ have isomorphic singularities and so $\text{dep}(Y) = \text{dep}(Y^+)$.

Suppose first that $f : X \rightarrow W$ is a flipping contraction. By induction, we then have $\text{dep}(Y) \geq \dots \geq \text{dep}(Y_i) \geq \dots \geq \text{dep}(Y')$. By induction on $\text{dep}(\dots)$ we obtain the required factorization for each $Y_i \dashrightarrow Y_{i+1}$ and hence for $X \dashrightarrow X'$.

Indeed, by Remark 3.4, either $Y' \rightarrow X'$ is a divisorial contraction to a curve, or at least one of $Y_i \rightarrow Y_{i+1}$ is a flip.

If $Y' \rightarrow X'$ is a divisorial contraction to a curve, then by induction, $\text{dep}(Y') \geq \text{dep}(X')$. Therefore,

$$\text{dep}(X) = \text{dep}(Y) + 1 > \text{dep}(Y) \geq \text{dep}(Y') \geq \text{dep}(X').$$

If $Y' \rightarrow X'$ is a divisorial contraction to a point, then by Remark 3.4, we have that at least one of $Y_i \rightarrow Y_{i+1}$ is a flip. Therefore, $\text{dep}(Y_i) > \text{dep}(Y_{i+1})$ by induction and hence $\text{dep}(Y) > \text{dep}(Y')$. Together with Proposition 2.15 that $\text{dep}(Y') > \text{dep}(X) - 1$, it follows that

$$\text{dep}(X) = \text{dep}(Y) + 1 > \text{dep}(Y') + 1 \geq \text{dep}(X').$$

Suppose next that $f : X \rightarrow W$ is a divisorial contraction to a curve. Again, we have $\text{dep}(Y) \geq \dots \geq \text{dep}(Y_i) \geq \dots \geq \text{dep}(Y')$. Since $Y' \rightarrow X'$ is a divisorial contraction to a curve and $\text{dep}(Y') < \text{dep}(X)$, we have $\text{dep}(Y') \geq \text{dep}(X')$. Finally, as $X' \rightarrow W$ is a divisorial contraction to a point, we have that $\text{dep}(X') \geq \text{dep}(W) - 1$. We thus conclude that

$$\text{dep}(X) = \text{dep}(Y) + 1 \geq \text{dep}(X') + 1 \geq \text{dep}(W).$$

Moreover, by induction on $\text{dep}(\dots)$ we obtain the required factorization for each $Y_i \dashrightarrow Y_{i+1}$ and hence for $X \dashrightarrow W$. $\square$

REFERENCES

- [1] C. Birkar, P. Cascini, C. Hacon, J. McKernan, *Existence of minimal models for varieties of log general type.*, J. Amer. Math. Soc., to appear. arXiv:math/0610203v2 [math.AG]
- [2] G. Brown, *Pluricanonical cohomology across flips*. Bull. London Math. Soc. **31** (1999), no. 5, 513–522.

- [3] J. A. Chen and M. Chen, *Explicit birational geometry of threefolds of general type, I*. Ann. Sci. École Norm. Sup., to appear. arXiv: 0810.5041.
- [4] A. Corti, *Singularities of linear systems and 3-fold birational geometry*. Explicit birational geometry of 3-folds, 259–312, London Math. Soc. Lecture Note Ser., 281, Cambridge Univ. Press, Cambridge, 2000.
- [5] S. D. Cutkosky, *Elementary contractions of Gorenstein threefolds*, Math. Ann. **280** (1988), no. 3, 521–525.
- [6] C. D. Hacon and J. McKernan, *Boundedness of pluricanonical maps of varieties of general type*, Invent. Math. **166** (2006), 1–25.
- [7] C. D. Hacon and J. McKernan, *Existence of minimal models for varieties of log general type II*. arXiv:0808.1929v1 [math.AG]
- [8] T. Hayakawa, *Blowing ups of 3-dimensional terminal singularities*, Publ. Res. Inst. Math. Sci. **35** (1999), no. 3, 515–570.
- [9] T. Hayakawa, *Blowing ups of 3-dimensional terminal singularities. II*, Publ. Res. Inst. Math. Sci. **36** (2000), no. 3, 423–456.
- [10] T. Hayakawa, *Gorenstein resolutions of 3-dimensional terminal singularities*, Nagoya Math. J. **178** (2005), 63–115.
- [11] T. Hayakawa, *Flips in dimension three via crepant descent method*. Proc. Japan Acad. Ser. A Math. Sci. **79** (2003), no. 2, 46–51.
- [12] T. Hayakawa, *Divisorial contractions to 3-dimensional terminal singularities with discrepancy one*. J. Math. Soc. Japan, **57** 2005.
- [13] Y. Kawamata, *Semistable minimal models of threefolds in positive or mixed characteristic*. J. Algebraic Geom. **3** (1994), no. 3, 463–491.
- [14] Y. Kawamata, *Divisorial contractions to 3-dimensional terminal quotient singularities*. Higher-dimensional complex varieties (Trento, 1994), 241–246, de Gruyter, Berlin, 1996.
- [15] Y. Kawamata, *Crepant blowing-up of 3-dimensional canonical singularities and its application to degenerations of surfaces*. Ann. of Math. (2) **127** (1988), no. 1, 93–163.
- [16] Y. Kawamata, *Flops connect minimal models*. (English summary) Publ. Res. Inst. Math. Sci. **44** (2008), no. 2, 419–423.
- [17] Y. Kawamata, K. Matsuda, K. Matsuki, *Introduction to the minimal model problem*, Algebraic geometry, Sendai, 1985, 283–360, Adv. Stud. Pure Math., **10**, North-Holland, Amsterdam, 1987.
- [18] M. Kawakita, *Divisorial contractions in dimension three which contract divisors to smooth points*, Invent. Math. **145** (2001), no. 1, 105–119.
- [19] M. Kawakita, *Divisorial contractions in dimension three which contract divisors to compound A_1 points*. Compositio Math. **133** (2002), no. 1, 95–116.
- [20] M. Kawakita, *General elephants of three-fold divisorial contractions*. J. Amer. Math. Soc. **16** (2003), no. 2, 331–362.
- [21] M. Kawakita, *Three-fold divisorial contractions to singularities of higher indices*, Duke Math. J. **130** (2005), no. 1, 57–126.
- [22] S. Keel, *Basepoint freeness for nef and big line bundles in positive characteristic*. Ann. of Math. (2) **149** (1999), no. 1, 253–286.
- [23] J. Kollár, *Flops*. Nagoya Math. J. **113** (1989), 15–36.
- [24] J. Kollár, *Extremal rays on smooth threefolds*. Ann. Sci. École Norm. Sup. (4) **24** (1991), no. 3, 339–361.
- [25] J. Kollár, *Singularities of pairs*. Algebraic geometry—Santa Cruz 1995, 221–287, Proc. Sympos. Pure Math., **62**, Part 1, Amer. Math. Soc., Providence, RI, 1997.
- [26] J. Kollár, *Higher direct images of dualizing sheaves I*, Ann. Math. **123**(1986), 11–42; II, *ibid.* **124**(1986), 171–202.

- [27] J. Kollár et. al, *Flips and abundance for algebraic threefolds*. Papers from the Second Summer Seminar on Algebraic Geometry held at the University of Utah, Salt Lake City, Utah, August 1991. Asterisque No. 211 (1992). Societe Mathematique de France, Paris, 1992. pp. 1–258.
- [28] H. Clemens, J. Kollár and S. Mori, *Higher-dimensional complex geometry*. Astérisque No. 166 (1988), 144 pp. (1989).
- [29] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, 1998, Cambridge Univ. Press.
- [30] J. Kollár and S. Mori, *Classification of three-dimensional flips*, J. Amer. Math. Soc. **5** (1992), no. 3, 533–703.
- [31] T. Luo, *Divisorial extremal contractions of threefolds: divisor to point*. Amer. J. Math. **120** (1998), no. 2, 441–451.
- [32] S. Mori, *Threefolds whose canonical bundles are not numerically effective*, Ann. Math. **116** (1982), 133–176.
- [33] S. Mori, *On 3-dimensional terminal singularities*, Nagoya Math. J. **98** (1985), 43–66.
- [34] S. Mori, *Flip theorem and the existence of minimal models for 3-folds*, J. Amer. Math. Soc. **1** (1988), no. 1, 117–253.
- [35] S. Mori, *Errata to: “Classification of three-dimensional flips” [J. Amer. Math. Soc. 5 (1992), no. 3, 533–703; MR1149195] by J. Kollár and Mori*. J. Amer. Math. Soc. **20** (2007), no. 1, 269–271
- [36] M. Reid, *Minimal models of canonical 3-folds*, Adv. Stud. Pure Math. **1**(1983), 131–180.
- [37] N. Tziolas, *Terminal 3-fold divisorial contractions of a surface to a curve. I*. Compositio Math. **139** (2003), no. 3, 239–261.
- [38] N. Tziolas, *Families of D-minimal models and applications to 3-fold divisorial contractions*. Proc. London Math. Soc. (3) **90** (2005), no. 2, 345–370.
- [39] N. Tziolas, *Three dimensional divisorial extremal neighborhoods*. Math. Ann. **333** (2005), no. 2, 315–354.
- [40] N. Tziolas, *$\mathbb{Q}$ -Gorenstein deformations of nonnormal surfaces*. Amer. J. Math. **131** (2009), no. 1, 171–193.
- [41] N. Tziolas, *3-fold divisorial extremal neighborhoods over $cE7$ and $cE6$ Compound DuVal singularities*. arXiv:math/0703855
- [42] M. Reid, *Young person’s guide to canonical singularities*, Proc. Symposia in pure Math. **46**(1987), 345–414.

DEPARTMENT OF MATHEMATICS, NATIONAL TAIWAN UNIVERSITY, TAIPEI,
106, TAIWAN

E-mail address: jkchen@math.ntu.edu.tw

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY,
155 SOUTH 1400 EAST, JWB 233, UT 84112-0090, USA

E-mail address: hacon@math.utah.edu